

UK Pesticide Guide

CLAYTON LANARK

Product Details



Registration Number:	12942	Marketing Company:	Clayton
Formulation:	Capsule suspension	Active Substance:	lambda-cyhalothrin
Pesticide Contents:	100 g/l	Pack Size:	1 litre
For use only as:	Agricultural insecticide, Horticultural insecticide	Transport Code:	9
Packaging Group:	3	UN Number:	3082
Mode of Action Code:	IRAC 3	Approval Expiry:	30 June 2024

Statutory Information Compliance with these conditions of use is a legal requirement

Targets	Maximum Dose	Maximum Treatments	Latest Application	Harvest Interval	Restrictions	* EAMU Number	EAMU Expiry
Broccoli Caterpillars, Whitefly	100 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Not stated	[No Data]	Min spray interval 10 d	[No Data]	[No Data]
Brussels sprouts Caterpillars, Whitefly	100 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Not stated	[No Data]	Min spray interval 10 d	[No Data]	[No Data]
Cabbages Caterpillars, Whitefly	100 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Not stated	[No Data]	Min spray interval 10 d	[No Data]	[No Data]
Calabrese Caterpillars, Whitefly	100 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Not stated	[No Data]	Min spray interval 10 d	[No Data]	[No Data]
Carrots Cutworms	75 ml/ha	150 ml/ha/crop (max 4 applications/crop)	[No Data]	14 days	Min spray interval 7 d	[No Data]	[No Data]
Cauliflowers Caterpillars, Whitefly	100 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Not stated	[No Data]	Min spray interval 10 d	[No Data]	[No Data]
Combining peas Pea and bean weevil, Pea aphid, Pea midge, Pea moth	75 ml/ha	150 ml/ha/crop (max 4 applications/crop)	Not stated	25 days	Min spray interval 7 d	[No Data]	[No Data]
Durum wheat Aphids, Barley yellow dwarf virus vectors	50 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Before late milk stage (GS 77)	[No Data]	Min spray interval 14 d	[No Data]	[No Data]
Edible podded peas Pea and bean weevil, Pea aphid, Pea midge, Pea moth	75 ml/ha	150 ml/ha/crop (max 4 applications/crop)	Not stated	[No Data]	Min spray interval 7 d	[No Data]	[No Data]
Lettuce Cutworms	75 ml/ha	150 ml/ha/crop (max 4 applications/crop)	[No Data]	7 days	Min spray interval 7 d	[No Data]	[No Data]
Pears Pear sucker	90 ml/ha	270 ml/ha/yr	Not stated	7 days	Min spray interval 14 d	[No Data]	[No Data]
Potatoes Aphids	75 ml/ha	300 ml/ha/crop (max 4 applications/crop)	Not stated	[No Data]	Min spray interval 7 d	[No Data]	[No Data]
Spring barley Aphids	50 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Before late milk stage (GS 77)	[No Data]	Min spray interval 14 d	[No Data]	[No Data]
Spring field beans	75 ml/ha	150 ml/ha/crop (max 4 applications/crop)	Not stated	25 days	Min spray interval 7 d	[No Data]	[No Data]

Targets	Maximum Dose	Maximum Treatments	Latest Application	Harvest Interval	Restrictions	* EAMU Number	EAMU Expiry
Pea and bean weevil		applications/crop)			d		
Spring oats Aphids	50 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Before watery ripe stage (GS 71)	[No Data]	Min spray interval 14 d	[No Data]	[No Data]
Spring oilseed rape Beet virus yellows vectors, Cabbage seed weevil, Cabbage stem flea beetle, Flea beetle, Pod midge, Pollen beetle	75 ml/ha	225 ml/ha/crop (max 4 applications/crop)	[No Data]	6 weeks	Min spray interval 7 d	[No Data]	[No Data]
Spring wheat Aphids	50 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Before late milk stage (GS 77)	[No Data]	Min spray interval 7 d	[No Data]	[No Data]
Sugar beet Beet leaf miner, Cutworms, Flea beetle	75 ml/ha	150 ml/ha/crop (max 4 applications/crop)	[No Data]	8 weeks	Min spray interval 7 d	[No Data]	[No Data]
Vining peas Pea and bean weevil, Pea aphid, Pea midge, Pea moth	75 ml/ha	150 ml/ha/crop (max 4 applications/crop)	Not specified	[No Data]	Min spray interval 7 d	[No Data]	[No Data]
Winter barley Aphids, Barley yellow dwarf virus vectors	50 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Before late milk stage (GS 77)	[No Data]	Min spray interval 7 d	[No Data]	[No Data]
Winter field beans Pea and bean weevil	75 ml/ha	150 ml/ha/crop (max 4 applications/crop)	Not stated	25 days	Min spray interval 7 d	[No Data]	[No Data]
Winter oats Aphids, Barley yellow dwarf virus vectors	50 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Before watery ripe stage (GS 71)	[No Data]	Min spray interval 7 d	[No Data]	[No Data]
Winter oilseed rape Beet virus yellows vectors, Cabbage seed weevil, Cabbage stem flea beetle, Flea beetle, Pod midge, Pollen beetle	75 ml/ha	225 ml/ha/crop (max 4 applications/crop)	Before end of flowering	[No Data]	Min spray interval 7 d	[No Data]	[No Data]
Winter wheat Aphids, Barley yellow dwarf virus vectors, Wheat-blossom midge, Yellow cereal fly	50 ml/ha	200 ml/ha/crop (max 4 applications/crop)	Before late milk stage (GS 77)	[No Data]	Min spray interval 7 d	[No Data]	[No Data]

Hazard classification & safety precautions

Hazard

- Dangerous for the environment
- Harmful

Operator protection

- Avoid contact with skin
- Wash all protective clothing thoroughly after use, especially the inside of gloves
- Wash concentrate/dust from skin or eyes immediately
- Wash hands and exposed skin before eating and drinking and after work
- When using do not eat, drink or smoke

Storage and disposal

- Keep away from food, drink and animal feeding-stuffs
- Keep in original container, tightly closed, in a safe place
- Keep out of reach of children
- Rinse container thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely
- This material (and its container) must be disposed of in a safe way

Risk phrases

- Harmful by inhalation
- Harmful if swallowed
- May cause long-term adverse effects in the aquatic environment
- May cause sensitization by skin contact
- Very toxic to aquatic organisms

Environmental protection

- Dangerous to bees
- Do not allow direct spray from broadcast air-assisted sprayers to fall within xx m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 5 m of the top of a ditch which is dry at the time of application. Aim spray away from water
- Do not allow direct spray from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water
- Do not re-use container for any purpose/Do not re-use container for any other purpose
- Use appropriate containment to avoid environmental contamination

Medical advice

- If you feel unwell, seek medical advice immediately (show the label where possible)

Efficacy

- Best results normally obtained from treatment when pest attack first seen. See label for detailed recommendations on each crop
- Timing for control of barley yellow dwarf virus vectors depends on specialist assessment of the level of risk in the area
- Repeat applications recommended in some crops where prolonged attack occurs, up to maximum total dose. See label for details
- Where strains of aphids resistant to lambda-cyhalothrin occur control is unlikely to be satisfactory
- Addition of wetter recommended for control of certain pests in brassicas and oilseed rape
- Use of sufficient water volume to ensure thorough crop penetration recommended for optimum results
- Use of drop-legged sprayer gives improved results in crops such as Brussels sprouts

Restrictions

- Maximum number of applications or maximum total dose per crop varies - see labels
- Do not apply to a cereal crop if any product containing a pyrethroid insecticide or dimethoate has been applied to the crop after the start of ear emergence (GS 51)
- Do not spray cereals in the spring/summer (i.e. after 1 Apr) within 6 m of edge of crop

Crop specific information

- Latest use before late milk stage on cereals; before end of flowering for winter oilseed rape HI 3 d for radishes, red beet; 7 d for lettuce [12629]; 14 d for carrots and parsnips; 25 d for peas, field beans; 6 wk for spring oilseed rape; 8 wk for sugar beet

Environmental safety

- Dangerous for the environment Very toxic to aquatic organisms Flammable [13457] To protect non-target arthropods respect an untreated buffer zone of 5 m to non-crop land Since there is a high risk to non-target insects or other arthropods, do not spray cereals in spring/summer i.e. after 1st April within 6m of the field boundary [18868]
Dangerous to bees